

Georgia Southern University Digital Commons@Georgia Southern

Mathematical Sciences Faculty Publications

Mathematical Sciences, Department of

7-7-2014

Existence of Positive Solutions for $p(x)$ -Laplacian Equations with a Singular Nonlinear Term

Jingjing Liu

Zhengzhou University of Light Industry

Qihu Zhang

Zhengzhou University of Light Industry

Chunshan Zhao

Georgia Southern University, czhao@georgiasouthern.edu

Follow this and additional works at: <https://digitalcommons.georgiasouthern.edu/math-sci-facpubs>

 Part of the [Mathematics Commons](#)

Recommended Citation

Liu, Jingjing, Qihu Zhang, Chunshan Zhao. 2014. "Existence of Positive Solutions for $p(x)$ -Laplacian Equations with a Singular Nonlinear Term." *Electronic Journal of Differential Equations*, 2014 (155): 1-21. source: <http://ejde.math.txstate.edu/Volumes/2014/155/abstr.html>

<https://digitalcommons.georgiasouthern.edu/math-sci-facpubs/323>

This article is brought to you for free and open access by the Mathematical Sciences, Department of at Digital Commons@Georgia Southern. It has been accepted for inclusion in Mathematical Sciences Faculty Publications by an authorized administrator of Digital Commons@Georgia Southern. For more information, please contact digitalcommons@georgiasouthern.edu.

EXISTENCE OF POSITIVE SOLUTIONS FOR $p(x)$ -LAPLACIAN EQUATIONS WITH A SINGULAR NONLINEAR TERM

JINGJING LIU, QIHU ZHANG, CHUNSHAN ZHAO

Dedicated to Professor Xianling Fan on his 70th birthday

ABSTRACT. In this article, we study the existence of positive solutions for the $p(x)$ -Laplacian Dirichlet problem

$$-\Delta_{p(x)}u = \lambda f(x, u)$$

in a bounded domain $\Omega \subset \mathbb{R}^N$. The singular nonlinearity term f is allowed to be either $f(x, s) \rightarrow +\infty$, or $f(x, s) \rightarrow +\infty$ as $s \rightarrow 0^+$ for each $x \in \Omega$. Our main results generalize the results in [15] from constant exponents to variable exponents. In particular, we give the asymptotic behavior of solutions of a simpler equation which is useful for finding supersolutions of differential equations with variable exponents, which is of independent interest.

1. INTRODUCTION

Let $\Omega \subset \mathbb{R}^N$ be an open bounded domain with C^2 boundary. We consider the existence of positive solutions for elliptic problems with variable exponent of the form

$$\begin{aligned} -\Delta_{p(x)}u &= \lambda f(x, u), & \text{in } \Omega, \\ u(x) &> 0, & \text{in } \Omega, \\ u(x) &= 0, & \text{on } \partial\Omega, \end{aligned} \tag{1.1}$$

where $-\Delta_{p(x)}u = -\operatorname{div}(|\nabla u|^{p(x)-2}\nabla u)$ with $\nabla u = (\partial_{x_1}u, \partial_{x_2}u, \dots, \partial_{x_N}u)$ which is so-called $p(x)$ -Laplacian, $p(\cdot)$ is a function which satisfies some conditions specified below, $f : \Omega \times (0, \infty) \rightarrow [0, \infty)$ is a continuous function, and $\lambda > 0$ is a real parameter. Throughout this paper, we will denote $d(x) = d(x, \partial\Omega)$.

In recent years, the study of differential equations and variational problems with nonstandard $p(x)$ -growth condition has been an interesting topic. The $p(x)$ -Laplacian arises from the study of nonlinear elasticity, electrorheological fluids and image restoration etc. For example, electrorheological fluids have an extensive applications in robotics, aircraft and aerospace. We refer readers to [1, 5, 19, 41, 42, 46] for more detailed background of applications. There are many reference papers related to the study of differential equations and variational problems with variable exponent. Far from being complete, we refer readers to [1, 2, 3, 6, 7, 8, 9, 10, 11,

2000 *Mathematics Subject Classification.* 35J25, 35J65, 35J70.

Key words and phrases. $p(x)$ -Laplacian; singular nonlinear term; sub-supersolution method.

©2014 Texas State University - San Marcos.

Submitted July 2, 2013. Published July 7, 2014.

12, 14, 20, 21, 22, 23, 24, 25, 26, 30, 31, 32, 33, 34, 35, 36, 39, 42, 43, 44, 45, 46] and references cited therein. For example, the regularity of weak solutions for differential equations with variable exponent was studied in [1, 7], and existence of solutions for variable exponent problems was studied in a series of papers [3, 9, 12, 20, 24, 30, 33, 36, 39, 44, 45]. Recently, the applications of variable exponent analysis in image restoration attracted more and more attention [16, 17, 23, 28]. In this paper, our aim is to study the existence of positive solution for problem (1.1) with singular nonlinear term f .

Clearly, if $p(\cdot) \equiv p$, a constant, the operator is the well-known p -Laplacian, and (1.1) is the usual p -Laplacian equation, but for non-constant $p(\cdot)$, $p(x)$ -Laplacian problems are more complicated due to the non-homogeneity of $p(x)$ -Laplacian. For example, if Ω is a smooth bounded domain, the Rayleigh quotient

$$\lambda_{p(\cdot)} = \inf_{u \in W_0^{1,p(\cdot)}(\Omega) \setminus \{0\}} \frac{\int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx}{\int_{\Omega} \frac{1}{p(x)} |u|^{p(x)} dx}$$

is zero in general, and $\lambda_{p(\cdot)} > 0$ only under some special conditions (see [13]). It is also possible the first eigenvalue and eigenfunction of $p(x)$ -Laplacian do not exist, even though the existence of the first eigenvalue and eigenfunction is very important in the study of elliptic problems related to p -Laplacian problems. For example, in [15], the author use the first eigenfunction and the first eigenvalue to construct subsolutions. Fan [8] considered the eigenvalue problem of $p(x)$ -Laplacian equation with the Neumann boundary condition, the existence of infinite many eigenvalues has been established. Benouhiba [2] studied the eigenvalue problem

$$-\Delta_{p(x)} u = \lambda V(x) |u|^{q(x)-2} u, \quad x \in \mathbb{R}^N,$$

where $1 < p(\cdot)$; $q(\cdot) \in C(\mathbb{R}^N)$ and $V(\cdot)$ is an indefinite weight function. The results show that the spectrum of such problems contains a continuous family of eigenvalues.

There are many papers deal with the existence of positive solution for a class p -Laplacian equation with singular nonlinearity (see [15, 18, 37, 38, 40] Mohammed [37], Perera and Silva —citep1, Qing and Yang [40] and Guo et al [18] studied the solvability of (1.1) with $\lambda = 1$, $p(\cdot) \equiv p \neq 2$ and $f(\cdot, \cdot)$ satisfies various conditions. In [38], the authors considered a boundary condition in a more general sense.

Mohammed [37] considered the existence and uniqueness of weak solutions of the singular boundary value problem with constant exponent as follows.

$$\begin{aligned} -\Delta_p u &= f(x, u), & \text{in } \Omega, \\ u(x) &> 0, & \text{in } \Omega, \\ u(x) &= 0, & \text{on } \partial\Omega, \end{aligned}$$

where Ω is a bounded domain in \mathbb{R}^N with $C^{1,\omega}$ boundary for some $0 < \omega < 1$, and singular nonlinearity term $f(x, t)$ could show up when $t \rightarrow 0^+$. Mohammed make the following two assumptions:

- (1) For each $\theta \in (0, 1)$, there is a constant $C_\theta \geq 1$ such that $g(\theta t) \leq C_\theta g(t)$ for all $t > 0$;
- (2) $f(x, s) \geq a(x)$ for any $(x, s) \in \Omega \times (0, \infty)$.

In [15], the authors studied the existence of solutions of the nonlinear elliptic problem with constant exponent,

$$\begin{aligned} -\Delta_p u &= \lambda f(x, u), \quad \text{in } \Omega, \\ u(x) &> 0, \quad \text{in } \Omega, \\ u(x) &= 0, \quad \text{on } \partial\Omega, \end{aligned}$$

where Ω is a bounded domain in \mathbb{R}^N , $1 < p < N$, $f : \Omega \times (0, \infty) \rightarrow [0, \infty)$ is a suitable function and $\lambda > 0$ is a real parameter. The nonlinearity term f is allowed to be either $f(x, s) \rightarrow +\infty$ or $f(x, s) \rightarrow +\infty$ as $s \rightarrow 0^+$ for each $x \in \Omega$, and the assumptions (1) and (2) are not assumed.

Results on elliptic problems with singular nonlinearity are rare (see [29, 44]). In [44], by using the sub-supersolution method, we studied the existence and the boundary asymptotic behavior of solutions of the elliptic problem with variable exponent,

$$\begin{aligned} -\Delta_{p(x)} u &= \frac{\lambda}{u^{\gamma(x)}}, \quad \text{in } \Omega, \\ u(x) &> 0, \quad \text{in } \Omega, \\ u(x) &= 0, \quad \text{on } \partial\Omega, \end{aligned}$$

where $\Omega \subset \mathbb{R}^N$ is a domain with C^2 boundary, λ is a positive parameter which is large enough.

Liu [29] generalized the results of [37] to $p(x)$ -Laplacian by making the similar assumptions. The condition (1) implies that $g(t) \leq Ct^{-a}$ when $t \leq 1$ for some $a > 0$, which is invalid for $g(t) = e^{1/t}$, and the condition (2) is a bit strong in some sense. Motivated by [15], in this article we partly generalized the results to $p(x)$ -Laplacian.

Before stating our main results, we make the following assumptions throughout this paper:

- (H0) $p(\cdot) \in C^1(\overline{\Omega})$, $1 < p^- := \inf_{\Omega} p(x) \leq p^+ := \sup_{\Omega} p(x) < \infty$, $p(\cdot) < N$.
- (H1) $f(x, s) \leq b(x)g(s)$ for all $(x, s) \in \Omega \times (0, \infty)$, where $g : (0, +\infty) \rightarrow (0, +\infty)$ is a continuous function, $sg(s)$ is decreasing for $s \leq 1$; and $b : \Omega \rightarrow [1, \infty)$, $b(\cdot) \in L^{\alpha(\cdot)}(\Omega)$, $1 < \alpha(\cdot) \in C(\overline{\Omega})$, and $\frac{1}{\alpha(x)} + \frac{1}{p^*(x)} < 1$, for all $x \in \overline{\Omega}$.
- (H2) $f(x, s)$ satisfies

$$\liminf_{s \rightarrow 0^+} \frac{f(x, s)}{s^{p^- - 1} |\ln s|^{p^-}} = +\infty \quad \text{uniformly for } x \in \Omega. \quad (1.2)$$

Theorem 1.1. *Assume that (H0), (H1), (H2) hold. Then problem (1.1) has a solution when λ is small enough.*

Theorem 1.2. *Assume that (H0), (H1), (H2) hold. Also assume that*

- (i) *there is a small $\delta > 0$ such that $p(x) \equiv p$ (a constant) for any $x \in \Omega$ with $d(x) \leq \delta$;*
- (ii) $\lim_{s \rightarrow +\infty} \frac{g(s)}{s^{p^- - 1 - \varepsilon}} := g_\infty \in [0, +\infty)$, *where $\varepsilon > 0$ is small enough;*
- (iii) $\alpha(\cdot) > N$ *on $\overline{\Omega}$.*

Then problem (1.1) has a solution for any positive λ .

Theorem 1.3. *Assume that (H0), (H1), (H2) hold. Also assume that*

- (i) $\frac{\partial p(\cdot)}{\partial \nu} < 0$ *on $\partial\Omega$, where ν is the inward unit normal vector of $\partial\Omega$;*

- (ii) $\lim_{s \rightarrow +\infty} \frac{g(s)}{s^{p^*-1-\varepsilon}} := g_\infty \in [0, +\infty)$, where $\varepsilon > 0$ is small enough;
 (iii) $\alpha(\cdot) > N$ on $\bar{\Omega}$.

Then problem (1.1) has a solution for any positive constant λ .

Theorem 1.4. Assume that (H0), (H1), (H2) hold. Also assume that

- (i) Equation (1.1) is radial;
 (ii) $\lim_{s \rightarrow +\infty} \frac{g(s)}{s^{p^*-1-\varepsilon}} := g_\infty \in [0, +\infty)$, where $\varepsilon > 0$ is small enough.

Then problem (1.1) has a solution for any positive λ .

This paper is organized as follows. In section 2, we will recall some basic facts about the variable exponent Lebesgue and Sobolev spaces which we will use later, and we will also give a general principle of sub-supersolution method. Proofs of our results will be presented in section 3.

2. PRELIMINARIES

Throughout this paper, the letters c, c_i, C, C_i ($i = 1, 2, \dots$), denote positive constants which may vary from line to line, but they are independent of the terms which will take part in any limit process.

To deal with the $p(x)$ -Laplacian problem, we need introduce some functional spaces $L^{p(\cdot)}(\Omega)$, $W^{1,p(\cdot)}(\Omega)$, $W_0^{1,p(\cdot)}(\Omega)$ and properties of the $p(x)$ -Laplacian which we will use later. Denote by $S(\Omega)$ be the set of all measurable real-valued functions defined in Ω . Note that two measurable functions are considered as the same element of $S(\Omega)$ when they are equal almost everywhere. Let

$$L^{p(\cdot)}(\Omega) = \left\{ u \in S(\Omega) : \int_{\Omega} |u(x)|^{p(x)} dx < \infty \right\},$$

with the norm

$$|u|_{p(\cdot)} = |u|_{L^{p(\cdot)}(\Omega)} = \inf \left\{ \lambda > 0 : \int_{\Omega} \left| \frac{u(x)}{\lambda} \right|^{p(x)} dx \leq 1 \right\}.$$

The space $(L^{p(\cdot)}(\Omega), |\cdot|_{p(\cdot)})$ becomes a Banach space. We call it variable exponent Lebesgue space. Moreover, this space is a separable, reflexive and uniform convex Banach space; see [14, Theorems 1.6, 1.10, 1.14].

The variable exponent Sobolev space

$$W^{1,p(\cdot)}(\Omega) = \left\{ u \in L^{p(\cdot)}(\Omega) : |\nabla u| \in L^{p(\cdot)}(\Omega) \right\},$$

can be equipped with the norm

$$\|u\| = |u|_{p(\cdot)} + |\nabla u|_{p(\cdot)}, \quad \forall u \in W^{1,p(\cdot)}(\Omega).$$

Note that $W_0^{1,p(\cdot)}(\Omega)$ is the closure of $C_0^\infty(\Omega)$ in $W^{1,p(\cdot)}(\Omega)$. The spaces $W^{1,p(\cdot)}(\Omega)$ and $W_0^{1,p(\cdot)}(\Omega)$ are separable, reflexive and uniform convex Banach spaces (see [14, Theorem 2.1]).

For $u, v \in S(\Omega)$, we write $u \leq v$ if $u(x) \leq v(x)$ for a.e. $x \in \Omega$. Let $\rho(x, s)$ be a Carathéodory function on $\Omega \times \mathbb{R}$ with property that for any $s_0 > 0$ there exists a constant A such that

$$|\rho(x, s)| \leq A \quad \text{for a.e. } x \in \Omega \text{ and all } s \in [-s_0, s_0]. \quad (2.1)$$

Definition 2.1. (i) Let $\underline{u}, \bar{u} \in W_{\text{loc}}^{1,p(\cdot)}(\Omega) \cap C_0(\bar{\Omega})$ satisfy $\underline{u}, \bar{u} > 0$ in Ω . We say \underline{u} and \bar{u} are a subsolution and a supersolution of (1.1) respectively, if

$$\begin{aligned} \int_{\Omega} |\nabla \underline{u}|^{p(x)-2} \nabla \underline{u} \nabla \phi \, dx &\leq \int_{\Omega} \lambda f(x, \underline{u}) \phi \, dx, \\ \int_{\Omega} |\nabla \bar{u}|^{p(x)-2} \nabla \bar{u} \nabla \phi \, dx &\geq \int_{\Omega} \lambda f(x, \bar{u}) \phi \, dx, \end{aligned}$$

for all $\phi \in C_0^\infty(\Omega)$ with $\phi \geq 0$ and $\text{supp } \phi \subset \subset \Omega$. We say u is a solution of (1.1), if it is both a subsolution and a supersolution of (1.1).

(ii) A function $u \in W^{1,p(\cdot)}(\Omega) \cap C(\bar{\Omega})$ is called a weak solution of the problem

$$\begin{aligned} -\Delta_{p(x)} u &= \rho(x, u), \quad \text{in } \Omega, \\ u(x) &= \varphi(x), \quad \text{on } \Omega, \end{aligned} \tag{2.2}$$

where $\varphi(\cdot) \in C(\bar{\Omega})$, if

$$\int_{\Omega} |\nabla u|^{p(x)-2} \nabla u \nabla \phi \, dx = \int_{\Omega} \rho(x, u) \phi \, dx, \quad \forall \phi \in C_0^\infty(\Omega);$$

(iii) $\underline{u}, \bar{u} \in W^{1,p(\cdot)}(\Omega) \cap C(\bar{\Omega})$ are called a weak subsolution and a weak supersolution of the problem (2.2) respectively if $\underline{u} \leq \varphi$ and $\bar{u} \geq \varphi$ on $\partial\Omega$ and for all $\phi \in C_0^\infty(\Omega)$, $\phi \geq 0$,

$$\begin{aligned} \int_{\Omega} |\nabla \underline{u}|^{p(x)-2} \nabla \underline{u} \nabla \phi \, dx &\leq \int_{\Omega} \rho(x, \underline{u}) \phi \, dx, \\ \int_{\Omega} |\nabla \bar{u}|^{p(x)-2} \nabla \bar{u} \nabla \phi \, dx &\geq \int_{\Omega} \rho(x, \bar{u}) \phi \, dx. \end{aligned}$$

Lemma 2.2 ([12, Proposition 2.1]). *The space $(L^{p(\cdot)}(\Omega), |\cdot|_{p(\cdot)})$ is a separable, uniform convex Banach space, and its conjugate space is $L^{p^0(\cdot)}(\Omega)$, where $p^0(\cdot)$ is the conjugate function of $p(\cdot)$ satisfying $\frac{1}{p(\cdot)} + \frac{1}{p^0(\cdot)} \equiv 1$. For any $u \in L^{p(\cdot)}(\Omega)$ and $v \in L^{p^0(\cdot)}(\Omega)$, we have the following Hölder inequality*

$$\left| \int_{\Omega} uv \, dx \right| \leq \int_{\Omega} |uv| \, dx \leq \left(\frac{1}{p^-} + \frac{1}{(p^0)^-} \right) |u|_{p(\cdot)} |v|_{p^0(\cdot)} \leq 2 |u|_{p(\cdot)} |v|_{p^0(\cdot)}.$$

Definition 2.3. Let $u, v \in W^{1,p(\cdot)}(\Omega) \cap L^\infty(\Omega)$. We say that $-\Delta_{p(x)} u + \rho(x, u) \leq -\Delta_{p(x)} v + \rho(x, v)$ in Ω if

$$\int_{\Omega} |\nabla u|^{p(x)-2} \nabla u \nabla \phi \, dx + \int_{\Omega} \rho(x, u) \phi \, dx \leq \int_{\Omega} |\nabla v|^{p(x)-2} \nabla v \nabla \phi \, dx + \int_{\Omega} \rho(x, v) \phi \, dx$$

for all $\phi \in C_0^\infty(\Omega)$, $\phi \geq 0$.

Next we give a comparison principle as follows.

Lemma 2.4 ([43, Lemma 2.3]). *Let $\rho(x, t)$ be a function satisfying (2.1) and non-decreasing in t . Let $u, v \in W^{1,p(\cdot)}(\Omega)$ satisfy*

$$-\Delta_{p(x)} u + \rho(x, u) \leq -\Delta_{p(x)} v + \rho(x, v), \quad (x \in \Omega),$$

if $u \leq v$ on $\partial\Omega$, then $u \leq v$ in Ω .

Lemma 2.5 ([6, Theorem 8.3.1]). *For every $u \in W_0^{1,p(\cdot)}(\Omega)$, the inequality*

$$|u|_{p^*(\cdot)} \leq C |\nabla u|_{p(\cdot)}$$

holds with a constant C depending only on the dimension N and p^+ and independent of Ω .

Lemma 2.6. *Suppose the domain Ω has finite measure, i.e. $|\Omega| < +\infty$, $p(\cdot), q(\cdot) \in C(\overline{\Omega})$, and $1 < p(x) < q(x) < N$, for all $x \in \overline{\Omega}$. Then for every $u \in L^{q(\cdot)}(\Omega)$, the following inequality holds:*

$$|u|_{p(\cdot)} \leq 2|\Omega|^{\frac{1}{p(\xi)} - \frac{1}{q(\xi)}} |u|_{q(\cdot)}, \quad \text{where } \xi \in \overline{\Omega}.$$

Moreover, $|u|_{p(\cdot)} \leq 2|\Omega|^{1/N} |u|_{p^*(\cdot)}$ for all $u \in W_0^{1,p(\cdot)}(\Omega)$.

The basic principle of sub-supersolution method for (1.1) can be stated as follows.

Lemma 2.7. *Suppose that (H0) holds and $\underline{u}, \bar{u} \in W_{\text{loc}}^{1,p(\cdot)}(\Omega) \cap C_0(\overline{\Omega})$. Let \underline{u} and \bar{u} be a subsolution and a supersolution of (1.1) respectively satisfying $\underline{u} \leq \bar{u}$. If $f \in C(\overline{\Omega} \times \mathbb{R}, \mathbb{R})$, then (1.1) has a solution $u \in W_{\text{loc}}^{1,p(\cdot)}(\Omega) \cap C_0(\overline{\Omega})$ satisfy $\underline{u} \leq u \leq \bar{u}$.*

Proof. Denote $\Omega_n = \{x \in \Omega : d(x) > 1/n\}$. Let

$$\tilde{f}(x, u) = \begin{cases} f(x, \bar{u}), & u \geq \bar{u}, \\ f(x, u), & \underline{u} < u < \bar{u}, \\ f(x, \underline{u}), & u \leq \underline{u}. \end{cases}$$

Consider

$$\begin{aligned} -\Delta_{p(x)} u &= \lambda \tilde{f}(x, u), \quad \text{in } \Omega_n, \\ u(x) &> 0, \quad \text{in } \Omega_n, \\ u(x) &= \underline{u}(x), \quad \text{on } \partial\Omega_n. \end{aligned} \tag{2.3}$$

Since $|\tilde{f}(x, u)|$ is bounded on $\overline{\Omega_n}$ and $\underline{u} \in C_0(\overline{\Omega})$, it is easy to see that (2.3) has a solution u_n , satisfy $\underline{u} \leq u_n \leq \bar{u}$. By [11, Theorem 1.2], we can see that $\{u_n\}_{n \geq n_0+1}$ has uniformly bounded $C^{1,\alpha}$ norm on $\overline{\Omega_{n_0}}$. By the diagonal method, we can choose a subsequence $\{u_{n_k}\}$ of $\{u_n\}$ such that

$$u_{n_k}(x) \rightarrow u(x), \quad \nabla u_{n_k}(x) \rightarrow \nabla u(x), \quad \forall x \in \Omega,$$

where $u \in C_0(\overline{\Omega}) \cap C^1(\Omega)$. Thus u is a solution of (1.1) and satisfies $\underline{u} \leq u \leq \bar{u}$. \square

3. PROOFS OF MAIN RESULTS

To study the existence of solutions of (1.1), we need to do some preparation work. Note that by [10, Theorem 4.2], the following problem has a weak solution $\omega_b \in W_0^{1,p(\cdot)}(\Omega)$,

$$\begin{aligned} -\Delta_{p(x)} \omega &= b(x), \quad \text{in } \Omega, \\ \omega(x) &= 0, \quad \text{on } \partial\Omega. \end{aligned} \tag{3.1}$$

Since $b(\cdot)$ is nonnegative, by the comparison principle it follows that ω_b is non-negative (see [43, Lemma 2.3]) and it is positive in Ω (see [43, Theorem 1.1]). From [10, Theorem 4.1], we see that ω_b is bounded. Then we have $\omega_b \in C^{1,\alpha}(\overline{\Omega})$ and $\frac{\partial \omega_b}{\partial \nu} > 0$ on $\partial\Omega$ from the following Lemma.

Lemma 3.1. (i) [7, Theorem 1.2] *Let ω_b be a bounded solution of (3.1), then $\omega_b \in C^{1,\alpha}(\overline{\Omega})$;*

(ii) [43, Theorem 1.2] *Let ω_b be a solution of (3.1), $x_1 \in \partial\Omega$, $\omega_b \in C^1(\Omega \cup \{x_1\})$, $\omega_b(x_1) = 0$. If Ω satisfies the inward-ball condition at x_1 , then $\frac{\partial \omega_b}{\partial \nu}(x_1) > 0$, where ν is the inward unit normal vector of $\partial\Omega$ on x_1 .*

We will prove the Theorems 1.1–1.4 stated in section 1 by using Lemma 2.7. Next we will construct a supersolution of (1.1) when λ is small enough. Before we begin the proof of Theorem 1.1, we need some background.

Define

$$g_{\#}(s) = \begin{cases} g(s), & \text{when } s < 1, \\ g(s), & \text{when } s \geq 1 \text{ and } \limsup_{s \rightarrow +\infty} \frac{g(s)}{s^{p^- - 1 - \varepsilon}} < +\infty, \\ g(1)s^{p^- - 1 - \varepsilon}, & \text{when } s \geq 1 \text{ and } \limsup_{s \rightarrow +\infty} \frac{g(s)}{s^{p^- - 1 - \varepsilon}} = +\infty. \end{cases}$$

Without loss of generality, we assume that $g_{\#}(s) = C_* s^{p^- - 1 - \varepsilon}$ for $s \geq 1$. There exists $M_0 = M_0(\delta)$ large enough such that

$$g_{\#}(s) < \delta s^{p^- - 1}, \quad \forall s \geq M_0. \quad (3.2)$$

Now we define a continuous function $\hat{g} : (0, \infty) \rightarrow (0, \infty)$ by

$$\hat{g}(s) := \sup \left\{ \frac{g_{\#}(t)}{t^{p^- - 1}}, t > s \right\}, \quad s > 0.$$

It follows from (3.2) and the definition of \hat{g} that

- (i) \hat{g} is non-increasing;
- (ii) $\hat{g}(s) \geq \frac{g_{\#}(s)}{s^{p^- - 1}}, s > 0$;
- (iii) $\hat{g}(s) < \delta$, for all $s \geq M_0$.

We also define a C^1 -function

$$H(s) := \left(\frac{2}{s} \int_{\frac{s}{2}}^s \hat{g}(t) dt \right)^{\frac{1}{p^- - 1}}, \quad s > 0.$$

Lemma 3.2. *The function H satisfies*

- (i) H is strictly decreasing, and $-H'(s) \geq \frac{2^\varepsilon - 1}{p^- - 1} \frac{H(s)}{s}$;
- (ii) $\hat{g}(s) \leq [H(s)]^{p^- - 1} \leq \hat{g}(s/2)$, $s > 0$;
- (iii) $H(s) \rightarrow +\infty$ as $s \rightarrow 0^+$, $H(s) \rightarrow 0^+$, when $s \rightarrow +\infty$.

Proof. We only need to prove $-H'(s) \geq \frac{2^\varepsilon - 1}{p^- - 1} \frac{H(s)}{s}$, the rest is easy to be verified. By computations

$$-H'(s) = \frac{1}{p^- - 1} \left(\frac{2}{s} \int_{\frac{s}{2}}^s \hat{g}(t) dt \right)^{\frac{1}{p^- - 1} - 1} \left(\frac{2}{s^2} \int_{\frac{s}{2}}^s \hat{g}(t) dt + \frac{2}{s} \left(\frac{1}{2} \hat{g}\left(\frac{s}{2}\right) - \hat{g}(s) \right) \right).$$

By condition (H1), when $s \leq 1$, $s^{p^-} \hat{g}(s)$ is decreasing, then we have $\frac{1}{2} \hat{g}(\frac{s}{2}) - \hat{g}(s) \geq 0$, and then

$$-H'(s) \geq \frac{1}{p^- - 1} \left(\frac{2}{s} \int_{\frac{s}{2}}^s \hat{g}(t) dt \right)^{\frac{1}{p^- - 1} - 1} \frac{2}{s^2} \int_{\frac{s}{2}}^s \hat{g}(t) dt \geq \frac{2^\varepsilon - 1}{p^- - 1} \frac{H(s)}{s}.$$

Here we note that $\hat{g}(s) = C_* s^{-\varepsilon}$ for $s \geq 1$. When $s \geq 2$, we have

$$\begin{aligned} \frac{2}{s} \left(\frac{1}{2} \hat{g}\left(\frac{s}{2}\right) - \hat{g}(s) \right) &= C_* \frac{2}{s} \left(\frac{1}{2} \left(\frac{s}{2}\right)^{-\varepsilon} - s^{-\varepsilon} \right) \\ &= C_* \frac{2}{s} (2^{\varepsilon - 1} - 1) s^{-\varepsilon} \\ &= \frac{2}{s} (2^{\varepsilon - 1} - 1) \hat{g}(s) \\ &\geq \frac{2}{s} (2^{\varepsilon - 1} - 1) \frac{2}{s} \int_{\frac{s}{2}}^s \hat{g}(t) dt \end{aligned}$$

$$= (2^\varepsilon - 2) \frac{2}{s^2} \int_{\frac{s}{2}}^s \hat{g}(t) dt,$$

and

$$\begin{aligned} -H'(s) &= \frac{1}{p^- - 1} \left(\frac{2}{s} \int_{\frac{s}{2}}^s \hat{g}(t) dt \right)^{\frac{1}{p^- - 1} - 1} \left(\frac{2}{s^2} \int_{\frac{s}{2}}^s \hat{g}(t) dt + \frac{2}{s} \left(\frac{1}{2} \hat{g}\left(\frac{s}{2}\right) - \hat{g}(s) \right) \right) \\ &\geq \frac{1}{p^- - 1} \left(\frac{2}{s} \int_{\frac{s}{2}}^s \hat{g}(t) dt \right)^{\frac{1}{p^- - 1} - 1} \left(\frac{2}{s^2} \int_{\frac{s}{2}}^s \hat{g}(t) dt + (2^\varepsilon - 2) \frac{2}{s^2} \int_{\frac{s}{2}}^s \hat{g}(t) dt \right) \\ &= \frac{1}{p^- - 1} \left(\frac{2}{s} \int_{\frac{s}{2}}^s \hat{g}(t) dt \right)^{\frac{1}{p^- - 1} - 1} \left(\frac{2}{s^2} (2^\varepsilon - 1) \int_{\frac{s}{2}}^s \hat{g}(t) dt \right) \\ &= \frac{2^\varepsilon - 1}{p^- - 1} \frac{H(s)}{s}. \end{aligned}$$

Note that $s^{p^-} \hat{g}(s)$ is decreasing for $s \leq 1$. When $1 < s < 2$, we have

$$\begin{aligned} \frac{2}{s} \left(\frac{1}{2} \hat{g}\left(\frac{s}{2}\right) - \hat{g}(s) \right) &= \frac{2}{s} \left(\frac{1}{2} \left(\frac{s}{2}\right)^{-p^-} \left(\frac{s}{2}\right)^{p^-} \hat{g}\left(\frac{s}{2}\right) - C_* s^{-\varepsilon} \right) \\ &\geq \frac{2}{s} \left(\frac{1}{2} \left(\frac{s}{2}\right)^{-p^-} \left(\frac{2}{s}\right)^{p^-} \hat{g}\left(\frac{2}{s}\right) - C_* s^{-\varepsilon} \right) \\ &= C_* \frac{2}{s} \left(\frac{1}{2} \left(\frac{s}{2}\right)^{-p^-} - s^{-\varepsilon} \right) \\ &= C_* \frac{1}{s} s^{-\varepsilon} (2^{p^-} s^{\varepsilon - p^-} - 2) \\ &\geq C_* \frac{1}{s} s^{-\varepsilon} (2^\varepsilon - 2) \\ &\geq (2^\varepsilon - 2) \frac{2}{s^2} \int_{\frac{s}{2}}^s \hat{g}(t) dt. \end{aligned}$$

Then we have

$$\begin{aligned} -H'(s) &= \frac{1}{p^- - 1} \left(\frac{2}{s} \int_{\frac{s}{2}}^s \hat{g}(t) dt \right)^{\frac{1}{p^- - 1} - 1} \left(\frac{2}{s^2} \int_{\frac{s}{2}}^s \hat{g}(t) dt + \frac{2}{s} \left(\frac{1}{2} \hat{g}\left(\frac{s}{2}\right) - \hat{g}(s) \right) \right) \\ &\geq \frac{1}{p^- - 1} \left(\frac{2}{s} \int_{\frac{s}{2}}^s \hat{g}(t) dt \right)^{\frac{1}{p^- - 1} - 1} \left(\frac{2}{s^2} \int_{\frac{s}{2}}^s \hat{g}(t) dt + (2^\varepsilon - 2) \frac{2}{s^2} \int_{\frac{s}{2}}^s \hat{g}(t) dt \right) \\ &= \frac{1}{p^- - 1} \left(\frac{2}{s} \int_{\frac{s}{2}}^s \hat{g}(t) dt \right)^{\frac{1}{p^- - 1} - 1} \left(\frac{2}{s^2} (2^\varepsilon - 1) \int_{\frac{s}{2}}^s \hat{g}(t) dt \right) \\ &= \frac{2^\varepsilon - 1}{p^- - 1} \frac{H(s)}{s}. \end{aligned}$$

By summarizing the above discussion, we have

$$-H'(s) \geq \frac{2^\varepsilon - 1}{p^- - 1} \frac{H(s)}{s}, \quad \forall s > 0.$$

The proof is complete. \square

As a consequence of Lemma 3.2, we can define the function

$$\eta(s) := \int_0^s \frac{1}{H(t)} dt, \quad s \geq 0, \quad (3.3)$$

for it is easy to show that $\eta \in C^2(0, \infty)$.

Lemma 3.3. *The function η satisfies*

- (i) $\eta : (0, \infty) \rightarrow (0, \infty)$ is strictly increasing;
- (ii) let $\psi = \eta^{-1}$ be the inverse function of η . Then $\psi'(s) = H(\psi(s))$, $s > 0$.

Denote $\Omega_\sigma = \{x \in \Omega : \omega_b(x) < \sigma\}$, where $\sigma > 0$ is a small positive constant.

Lemma 3.4. *Assume that (H0) and (H1) hold. Then there is a supersolution v of (1.1) such that $v \in W_{\text{loc}}^{1,p(\cdot)}(\Omega) \cap C_0(\overline{\Omega})$ when λ is small enough.*

Proof. Define

$$v(x) := \begin{cases} \psi(k_1 \omega_b(x)), & x \in \Omega_\sigma, \\ \omega_b(x) + \psi(k_1 \sigma) - \sigma, & x \in \Omega \setminus \Omega_\sigma, \end{cases} \quad (3.4)$$

where ω_b is given by (3.1), and $k_1 > 1$ is a constant. Obviously, $v \in C_0(\overline{\Omega}) \cap W_{\text{loc}}^{1,p(\cdot)}(\Omega)$. From the definition of g and \hat{g} , Lemma 3.2, and Lemma 3.3, it follows that

$$\begin{aligned} \psi'(k_1 \omega_b(x)) &= H(\psi(k_1 \omega_b(x))) = H(v(x)), x \in \Omega_\sigma, \\ \psi''(s) &\leq 0, \text{ for all } s \geq 0. \end{aligned}$$

We will prove this Lemma in three steps.

Step 1. We will prove that v is a super-solution of (1.1) in Ω_σ ; i.e.,

$$\int_{\Omega} |\nabla v|^{p(x)-2} \nabla v \nabla \phi \, dx \geq \int_{\Omega} \lambda b g(v) \phi \, dx \geq \int_{\Omega} \lambda f(x, v) \phi \, dx,$$

for any $\phi \in C_0^\infty(\Omega_\sigma)$ with $\phi \geq 0$ and $\text{supp } \phi \subset \subset \Omega_\sigma$. By computation, we have

$$\begin{aligned} & \int_{\Omega} |\nabla v|^{p(x)-2} \nabla v \nabla \phi \, dx \\ &= \int_{\Omega} [k_1 \psi'(k_1 \omega_b)]^{p(x)-1} |\nabla \omega_b|^{p(x)-2} \nabla \omega_b \nabla \phi \, dx \\ &= \int_{\Omega} |\nabla \omega_b|^{p(x)-2} \nabla \omega_b \nabla \left\{ \phi [k_1 \psi'(k_1 \omega_b)]^{p(x)-1} \right\} \, dx \\ & \quad - \int_{\Omega} (k_1)^{p(x)} |\nabla \omega_b|^{p(x)} \phi (p(x) - 1) [\psi'(k_1 \omega_b)]^{p(x)-2} \psi''(k_1 \omega_b) \, dx \\ & \quad - \int_{\Omega} (k_1)^{p(x)-1} |\nabla \omega_b|^{p(x)-2} \nabla \omega_b \nabla p [\phi \psi'(k_1 \omega_b)^{p(x)-1}] \ln k_1 \psi'(k_1 \omega_b) \, dx. \end{aligned}$$

By Lemma 3.2, we have $-H'(v) \geq \frac{2^\varepsilon - 1}{p^- - 1} \frac{H(v)}{v}$ which implies

$$-\psi''(k_1 \omega_b) = -H'(v) \psi'(k_1 \omega_b) = -H'(v) H(v) \geq \frac{2^\varepsilon - 1}{p^- - 1} \frac{H(v)}{v} H(v).$$

Note that $0 < c_1 \leq |\nabla \omega_b| \leq c_2$ on $\overline{\Omega_\sigma}$. Let σ be small enough. We can see that v is small enough in Ω_σ , and $H(v)$ is large enough in Ω_σ . Then we have

$$|\nabla \omega_b| \frac{2^\varepsilon - 1}{p^- - 1} k_1 \frac{H(v)}{v} \geq c_1 \frac{2^\varepsilon - 1}{p^- - 1} k_1 \frac{H(v)}{v} \geq |\nabla p| \ln k_1 H(v).$$

By computations, for any $\phi \in C_0^\infty(\Omega)$ with $\phi \geq 0$, we have

$$- \int_{\Omega} (k_1)^{p(x)} |\nabla \omega_b|^{p(x)} \phi (p(x) - 1) [\psi'(k_1 \omega_b)]^{p(x)-2} \psi''(k_1 \omega_b) \, dx$$

$$\begin{aligned}
&= - \int_{\Omega} (k_1)^{p(x)} |\nabla \omega_b|^{p(x)} \phi(p(x) - 1) [H(\psi(k_1 \omega_b))]^{p(x)-2} \psi''(k_1 \omega_b) dx \\
&= - \int_{\Omega} (k_1)^{p(x)} |\nabla \omega_b|^{p(x)} \phi(p(x) - 1) [H(v)]^{p(x)-2} H'(v) \psi'(k_1 \omega_b) dx \\
&\geq \int_{\Omega} (k_1)^{p(x)} |\nabla \omega_b|^{p(x)} \phi(p(x) - 1) [H(v)]^{p(x)-1} \frac{2^\varepsilon - 1}{p^- - 1} \frac{H(v)}{v} dx \\
&\geq \int_{\Omega} (k_1)^{p(x)-1} |\nabla \omega_b|^{p(x)-1} |\nabla p| [\phi \psi'(k_1 \omega_b)^{p(x)-1}] |\ln k_1 H(v)| dx.
\end{aligned}$$

Here we note that $H(s) \rightarrow +\infty$ as $s \rightarrow 0^+$ and $v \in C_0(\overline{\Omega})$. Then $v(x) \leq 1$ in Ω_σ and $H(v(x)) \geq 1$ for any $x \in \Omega_\sigma$ when σ is small enough. Thus we have

$$\begin{aligned}
\int_{\Omega} |\nabla v|^{p(x)-2} \nabla v \nabla \phi dx &\geq \int_{\Omega} |\nabla \omega_b|^{p(x)-2} \nabla \omega_b \nabla \{\phi [k_1 \psi'(k_1 \omega_b)]^{p(x)-1}\} dx \\
&= \int_{\Omega} b [k_1 \psi'(k_1 \omega_b)]^{p(x)-1} \phi dx \\
&\geq \int_{\Omega} b [k_1 H(v)]^{p^- - 1} \phi dx \\
&\geq \int_{\Omega} b k_1^{p^- - 1} \hat{g}(v) \phi dx \\
&\geq \int_{\Omega} b k_1^{p^- - 1} \frac{g(v)}{v^{p^- - 1}} \phi dx \\
&\geq \int_{\Omega} b g(v) \phi dx,
\end{aligned}$$

for any $\phi \in C_0^\infty(\Omega_\sigma)$ with $\phi \geq 0$ and $\text{supp } \phi \subset \subset \Omega$.

Then for any $\phi \in C_0^\infty(\Omega_\sigma)$ with $\phi \geq 0$ and $\text{supp } \phi \subset \subset \Omega_\sigma$, we have

$$\int_{\Omega} |\nabla v|^{p(x)-2} \nabla v \nabla \phi dx \geq \int_{\Omega} \lambda b(x) g(v) \phi dx \geq \int_{\Omega} \lambda f(x, v) \phi dx. \quad (3.5)$$

Step 2. We will prove that v is a supersolution of (1.1) in $\Omega \setminus \overline{\Omega_\sigma}$; i.e.,

$$\int_{\Omega} |\nabla v|^{p(x)-2} \nabla v \nabla \phi dx \geq \int_{\Omega} \lambda f(x, v) \phi dx, \forall \phi \in C_0^\infty(\Omega \setminus \overline{\Omega_\sigma}), \phi \geq 0.$$

Let λ be small enough such that $\lambda g(v(x)) \leq 1$, for all $x \in \Omega \setminus \overline{\Omega_\sigma}$. For any $\phi \in C_0^\infty(\Omega \setminus \overline{\Omega_\sigma})$ with $\phi \geq 0$, we have

$$\int_{\Omega} |\nabla v|^{p(x)-2} \nabla v \nabla \phi dx = \int_{\Omega} b \phi dx \geq \int_{\Omega} \lambda b g(v) \phi dx \geq \int_{\Omega} \lambda f(x, v) \phi dx. \quad (3.6)$$

Step 3. We will prove that v is a super-solution of (1.1) in Ω ; i.e.,

$$\int_{\Omega} |\nabla v|^{p(x)-2} \nabla v \nabla \phi dx \geq \int_{\Omega} \lambda f(x, v) \phi dx,$$

for all $\phi \in C_0^\infty(\Omega)$ with $\phi \geq 0$ and $\text{supp } \phi \subset \subset \Omega$. Denote $d_1(x) = d(x, \partial(\Omega \setminus \Omega_\sigma))$, and

$$\xi_n(x) = \begin{cases} 2n(\frac{1}{n} - d_1(x)), & \frac{1}{2n} \leq d_1(x) \leq \frac{1}{n}, \\ 1, & 0 \leq d_1(x) < \frac{1}{2n}, \quad n \in \mathbb{N} \\ 0, & \text{other wise.} \end{cases}$$

For $\phi \in C_0^\infty(\Omega_\sigma)$, with $\phi \geq 0$ and $\text{supp } \phi \subset \subset \Omega$, we have $\phi = \phi_{1,n} + \phi_{2,n} + \phi_{3,n}$, where $\phi_{1,n} = \phi \chi_{\overline{\Omega_\sigma}}(1 - \xi_n) \in C_0^\infty(\Omega)$ satisfies $\text{supp } \phi_{1,n} \Subset \Omega_\sigma$, $\phi_{2,n} = \phi \xi_n \in C_0^\infty(\Omega)$, and $\phi_{3,n} = \phi \chi_{\Omega \setminus \overline{\Omega_\sigma}}(1 - \xi_n) \in C_0^\infty(\Omega \setminus \overline{\Omega_\sigma})$. Therefore,

$$\begin{aligned} & \int_{\Omega} |\nabla v|^{p(x)-2} \nabla v \nabla \phi \, dx \\ &= \lim_{n \rightarrow \infty} \left\{ \int_{\Omega} |\nabla v|^{p(x)-2} \nabla v \nabla \phi_{1,n} \, dx + \int_{\Omega} |\nabla v|^{p(x)-2} \nabla v \nabla \phi_{2,n} \, dx \right. \\ & \quad \left. + \int_{\Omega} |\nabla v|^{p(x)-2} \nabla v \nabla \phi_{3,n} \, dx \right\} \\ &\geq \lim_{n \rightarrow \infty} \left\{ \int_{\Omega} \lambda f(x, v) \phi_{1,n} \, dx + \int_{\Omega} \lambda f(x, v) \phi_{3,n} \, dx + \int_{\Omega} |\nabla v|^{p(x)-2} \nabla v \nabla \phi_{2,n} \, dx \right\} \\ &= \int_{\Omega_\sigma} \lambda f(x, v) \phi \, dx + \int_{\Omega \setminus \Omega_\sigma} \lambda f(x, v) \phi \, dx + \lim_{n \rightarrow \infty} \int_{\Omega} |\nabla v|^{p(x)-2} \nabla v \nabla \phi_{2,n} \, dx \\ &= \int_{\Omega} \lambda f(x, v) \phi \, dx + \lim_{n \rightarrow \infty} \int_{\Omega} \phi |\nabla v|^{p(x)-2} \nabla v \nabla \xi_n \, dx \\ &= \int_{\Omega} \lambda f(x, v) \phi \, dx + \int_{\partial(\Omega \setminus \Omega_\sigma)} \phi [(\psi'(k_1 \sigma) |\nabla k_1 \omega_b|)^{p(x)-1} - |\nabla \omega_b|^{p(x)-1}] \, dS. \end{aligned}$$

Here we note that $\psi'(k_1 \sigma) = H(k_1 \sigma) \rightarrow +\infty$ as $\sigma \rightarrow 0^+$. Thus

$$\int_{\partial(\Omega \setminus \Omega_\sigma)} \phi [(\psi'(k_1 \sigma) |\nabla k_1 \omega_b|)^{p(x)-1} - |\nabla \omega_b|^{p(x)-1}] \, dS \geq 0,$$

and then

$$\int_{\Omega} |\nabla v|^{p(x)-2} \nabla v \nabla \phi \, dx \geq \int_{\Omega} \lambda f(x, v) \phi \, dx,$$

for all $\phi \in C_0^\infty(\Omega)$ with $\phi \geq 0$ and $\text{supp } \phi \subset \subset \Omega$. It means that v is a super-solution of (1.1). The proof is complete. \square

Proof of Theorem 1.1. At first, we construct a subsolution for problem (1.1). Since $\partial\Omega$ is C^2 smooth, there exists a positive constant ℓ such that $d(\cdot) \in C^2(\overline{\partial\Omega_{3\ell}})$, and $|\nabla d(\cdot)| \equiv 1$, where $\partial\Omega_{3\ell} = \{x \in \overline{\Omega} : d(x) \leq 3\ell\}$.

Let $\sigma \in (0, \ell)$ be small enough. Denote

$$\phi(x) = \begin{cases} e^{kd(x)} - 1, & d(x) < \sigma, \\ e^{k\sigma} - 1 + \int_{\sigma}^{d(x)} k e^{k\sigma} \left(\frac{2\ell-t}{2\ell-\sigma}\right)^{\frac{2}{p-1}} dt, & \sigma \leq d(x) < 2\ell, \\ e^{k\sigma} - 1 + \int_{\sigma}^{2\ell} k e^{k\sigma} \left(\frac{2\ell-t}{2\ell-\sigma}\right)^{\frac{2}{p-1}} dt, & 2\ell \leq d(x), \end{cases}$$

where $k > 0$ is a parameter. It is easy to see that $\phi \in C_0^1(\overline{\Omega})$. By computations it follows that

$$-\Delta_{p(x)\mu}\phi = \begin{cases} -k(k\mu e^{kd(x)})^{p(x)-1} [(p(x)-1) + (d(x) + \frac{\ln k\mu}{k}) \nabla p(x) \nabla d(x) + \frac{\Delta d(x)}{k}], & \text{if } d(x) < \sigma, \\ \left\{ \frac{1}{2\ell-\sigma} \frac{2(p(x)-1)}{p-1} - \left(\frac{2\ell-d(x)}{2\ell-\sigma}\right) [(\ln k\mu e^{k\sigma} \left(\frac{2\ell-d(x)}{2\ell-\sigma}\right)^{\frac{2}{p-1}}) \nabla p(x) \nabla d(x) + \Delta d(x)] \right\} (k\mu e^{k\sigma})^{p(x)-1} \left(\frac{2\ell-d(x)}{2\ell-\sigma}\right)^{\frac{2(p(x)-1)}{p-1}-1}, & \text{if } \sigma < d(x) < 2\ell, \\ 0 & \text{if } 2\ell < d(x). \end{cases}$$

Denote $\sigma = \frac{1}{k} \ln 2^{\frac{1}{p^+}}$. Then

$$e^{k\sigma} = 2^{\frac{1}{p^+}}. \quad (3.7)$$

Let

$$\mu = e^{-ak}, \quad \text{where } a = \frac{p^- - 1}{\max_{x \in \bar{\Omega}} |\nabla p| + 1}. \quad (3.8)$$

Then $k\mu \leq 1$ when k is large enough. If k is sufficiently large, it is easy to see that

$$-\Delta_{p(x)}\phi \leq 0 \leq \lambda f(x, \phi), d(x) < \sigma. \quad (3.9)$$

Since $d(\cdot) \in C^2(\bar{\Omega}_{3\ell})$, there exists a positive constant C_3 such that $|\Delta d| \leq C_3$ on $\bar{\Omega}_{3\ell}$. Note that the definition of σ means (3.7); i.e., $e^{k\sigma} = 2^{\frac{1}{p^+}}$. When k is large enough, we have

$$\begin{aligned} & -\Delta_{p(x)}\mu\phi \\ & \leq (k\mu e^{k\sigma})^{p(x)-1} \left(\frac{2\ell - d(x)}{2\ell - \sigma} \right)^{\frac{2(p(x)-1)}{p^- - 1} - 1} \left| \frac{2(p(x) - 1)}{(2\ell - \sigma)(p^- - 1)} \right. \\ & \quad \left. - \left(\frac{2\ell - d(x)}{2\ell - \sigma} \right) \left[(\ln k\mu e^{k\sigma}) \left(\frac{2\ell - d(x)}{2\ell - \sigma} \right)^{\frac{2}{p^- - 1}} \nabla p(x) \nabla d(x) + \Delta d(x) \right] \right| \\ & \leq C_1 (k\mu)^{p(x)-1} |\ln k + \ln \mu|, \quad \sigma < d(x) < 2\ell. \end{aligned} \quad (3.10)$$

When k is large enough, we can see that μ is small enough and moreover, $k\mu \leq 1$. Combining (1.2), (3.8) and (3.10) together, we have

$$\begin{aligned} -\Delta_{p(x)}\mu\phi & \leq C_1 (k\mu)^{p(x)-1} |\ln \mu| \\ & \leq C_1 (k\mu)^{p^- - 1} |\ln \mu| \\ & \leq C_1 a^{1-p^-} \mu^{p^- - 1} |\ln \mu|^{p^-} \\ & \leq C_2 (\mu\phi)^{p^- - 1} |\ln \mu\phi|^{p^-} \\ & \leq \lambda f(x, \mu\phi), \sigma < d(x) < 2\ell. \end{aligned} \quad (3.11)$$

Obviously

$$-\Delta_{p(x)}\mu\phi = 0 \leq \lambda f(x, \mu\phi), 2\ell < d(x). \quad (3.12)$$

Combining (3.9), (3.11) and (3.12), we can conclude that

$$-\Delta_{p(x)}\mu\phi \leq \lambda f(x, \mu\phi), \quad \text{a.e. in } \Omega. \quad (3.13)$$

Here we note that $\mu\phi = v$ on $\partial\Omega$, and $\mu\nabla\phi \leq \nabla v$ near $\partial\Omega$, then $\mu\phi \leq v$ on Ω when μ is small enough. By Lemma 2.7 and Lemma 3.4, (1.1) has a solution. The proof is complete. \square

To prove Theorem 1.2, we need the following Lemma, which is useful for finding supersolutions of (1.1). We denote by C_0 the best embedding constant of $W_0^{1,1}(\Omega) \subset L^{\frac{N}{N-1}}(\Omega)$ (see [4, Lemma 5.2]); i. e.,

$$|u|_{L^{N/(N-1)}(\Omega)} \leq C_0 |\nabla u|_{L^1(\Omega)} \quad \text{for } u \in W_0^{1,1}(\Omega). \quad (3.14)$$

Lemma 3.5. Suppose $0 \leq b(\cdot) \in L^{\alpha(\cdot)}(\Omega)$, $N < \alpha(\cdot) \in C(\bar{\Omega})$. Let $M > 0$ and u be the unique solution of the problem

$$\begin{aligned} -\operatorname{div}(|\nabla u|^{p(x)-2} \nabla u) &= Mb(x), \quad \text{in } \Omega, \\ u &= 0, \quad \text{on } \partial\Omega. \end{aligned} \quad (3.15)$$

Set

$$\rho_0 = \frac{p^-}{2|b(\cdot)|_{L^{\alpha^-}(\Omega)}|\Omega|^{\frac{1}{N}-\frac{1}{\alpha^-}}C_0}.$$

Then $|u|_\infty \leq C^* M^{1/(p^- - 1)}$ when $M \geq \rho_0$ and $|u|_\infty \leq C_* M^{1/(p^+ - 1)}$ when $M < \rho_0$, where C^* and C_* are positive constants depending only on $p^+, p^-, N, |b(\cdot)|_{L^{\alpha^-}(\Omega)}, |\Omega|$ and C_0 .

Proof. Let u be the solution of (3.15). Then $u \geq 0$. For $k \geq 0$, set $A_k = \{x \in \Omega : u(x) > k\}$. By taking $(u - k)^+$ as a test function of (3.15), it follows from (3.14) and Young inequality that

$$\begin{aligned} \int_{A_k} |\nabla u|^{p(x)} dx &= M \int_{A_k} b(u - k) dx \\ &\leq M |b(\cdot)|_{L^N(A_k)} |(u - k)^+|_{L^{N/(N-1)}(A_k)} \\ &\leq M |b(\cdot)|_{L^{\alpha^-}(A_k)} |A_k|^{\frac{1}{N}-\frac{1}{\alpha^-}} |(u - k)^+|_{L^{N/(N-1)}(A_k)} \\ &\leq M |b(\cdot)|_{L^{\alpha^-}(\Omega)} |A_k|^{\frac{1}{N}-\frac{1}{\alpha^-}} C_0 \int_{A_k} \varepsilon |\nabla u| \varepsilon^{-1} dx \\ &\leq M |b(\cdot)|_{L^{\alpha^-}(\Omega)} |A_k|^{\frac{1}{N}-\frac{1}{\alpha^-}} C_0 \int_{A_k} \left(\frac{(\varepsilon |\nabla u|)^{p(x)}}{p(x)} + \frac{(\varepsilon^{-1})^{p^0(x)}}{p^0(x)} \right) dx \\ &\leq \frac{M |b(\cdot)|_{L^{\alpha^-}(\Omega)} |\Omega|^{\frac{1}{N}-\frac{1}{\alpha^-}} C_0}{p^-} \int_{A_k} \varepsilon^{p(x)} |\nabla u|^{p(x)} dx \\ &\quad + \frac{M |b(\cdot)|_{L^{\alpha^-}(\Omega)} |A_k|^{\frac{1}{N}-\frac{1}{\alpha^-}} C_0}{(p^+)^0} \int_{A_k} \varepsilon^{-p^0(x)} dx. \end{aligned} \tag{3.16}$$

When $M \geq \rho_0$ we can take

$$\varepsilon = \left(\frac{p^-}{2M |b(\cdot)|_{L^{\alpha^-}(\Omega)} |\Omega|^{\frac{1}{N}-\frac{1}{\alpha^-}} C_0} \right)^{1/p^-} = \left(\frac{\rho_0}{M} \right)^{1/p^-}, \tag{3.17}$$

then $\varepsilon \leq 1$ and

$$\begin{aligned} &\frac{M |b(\cdot)|_{L^{\alpha^-}(\Omega)} |\Omega|^{\frac{1}{N}-\frac{1}{\alpha^-}} C_0}{p^-} \int_{A_k} \varepsilon^{p(x)} |\nabla u|^{p(x)} dx \\ &\leq \frac{M |b(\cdot)|_{L^{\alpha^-}(\Omega)} |\Omega|^{\frac{1}{N}-\frac{1}{\alpha^-}} C_0}{p^-} \varepsilon^{p^-} \int_{A_k} |\nabla u|^{p(x)} dx \\ &= \frac{1}{2} \int_{A_k} |\nabla u|^{p(x)} dx. \end{aligned}$$

Consequently, from the inequality above and (3.16) it follows that

$$\begin{aligned} \int_{A_k} |\nabla u|^{p(x)} dx &\leq \frac{2M |b(\cdot)|_{L^{\alpha^-}(\Omega)} |A_k|^{\frac{1}{N}-\frac{1}{\alpha^-}} C_0}{(p^+)^0} \int_{A_k} \varepsilon^{-p^0(x)} dx \\ &\leq \frac{2M |b(\cdot)|_{L^{\alpha^-}(\Omega)} C_0 \varepsilon^{-(p^-)^0}}{(p^+)^0} |A_k|^{1+\frac{1}{N}-\frac{1}{\alpha^-}}. \end{aligned} \tag{3.18}$$

Note that $b(\cdot) \geq 1$. From (3.15) and (3.18), we have

$$\int_{A_k} (u - k) dx \leq \int_{A_k} b(x)(u - k) dx = \frac{1}{M} \int_{A_k} |\nabla u|^{p(x)} dx \leq \gamma |A_k|^{1 + \frac{1}{N} - \frac{1}{\alpha^-}}, \quad (3.19)$$

where

$$\gamma = \frac{2|b(\cdot)|_{L^{\alpha^-}(\Omega)} C_0 \varepsilon^{-(p^-)^0}}{(p^+)^0}. \quad (3.20)$$

By the [27, Lemma 5.1, Chapter 2], (3.19) implies

$$|u|_\infty \leq \gamma \left(\frac{\alpha^- N}{\alpha^- - N} + 1 \right)^2 |\Omega|^{\frac{1}{N} - \frac{1}{\alpha^-}}. \quad (3.21)$$

From (3.17), (3.20) and (3.21), we see that

$$|u|_\infty \leq C^* M^{1/(p^- - 1)}, \quad (3.22)$$

where

$$C^* = \frac{(\frac{\alpha^- N}{\alpha^- - N} + 1)^2 (2C_0 |b(\cdot)|_{L^{\alpha^-}(\Omega)} |\Omega|^{\frac{1}{N} - \frac{1}{\alpha^-}})^{(p^-)^0}}{(p^+)^0 (p^-)^{(p^-)^0/p^-}}. \quad (3.23)$$

When $M < \rho_0$, take

$$\varepsilon = \left(\frac{p^-}{2M |b(\cdot)|_{L^{\alpha^-}(\Omega)} |\Omega|^{\frac{1}{N} - \frac{1}{\alpha^-}} C_0} \right)^{1/p^+} = \left(\frac{\rho_0}{M} \right)^{1/p^+}.$$

Note that in this case $\varepsilon > 1$. Using similar arguments as above we obtain

$$|u|_\infty \leq C_* M^{1/(p^+ - 1)},$$

where

$$C_* = \frac{(\frac{\alpha^- N}{\alpha^- - N} + 1)^2 (2C_0 |b(\cdot)|_{L^{\alpha^-}(\Omega)} |\Omega|^{\frac{1}{N} - \frac{1}{\alpha^-}})^{(p^+)^0}}{(p^+)^0 (p^-)^{(p^+)^0/p^+}}.$$

The proof is complete. \square

Lemma 3.6. Suppose there is a small $\delta > 0$ such that $p(x) \equiv p$ (a constant) for any $x \in \Omega$ with $d(x) \leq \delta$ and $N < \alpha(\cdot) \in C(\overline{\Omega})$. Let $M > 1$ and u be the unique solution of the problem

$$\begin{aligned} -\operatorname{div}(|\nabla u|^{p(x)-2} \nabla u) &= Mb(x), \quad \text{in } \Omega, \\ u &= 0, \quad \text{on } \partial\Omega, \end{aligned} \quad (3.24)$$

where $0 \leq b(\cdot) \in L^{\alpha(\cdot)}(\Omega)$. Then $|\nabla u(\cdot)| \leq CM^{\frac{1}{p^- - 1}}$ on $\partial\Omega$.

Proof. By Lemma 3.5, we have $u(x) \leq C_\# M^{\frac{1}{p^- - 1}}$ for all $x \in \Omega$. Let u_2 be the solution of the following p -Laplacian equation (with constant exponent)

$$\begin{aligned} -\operatorname{div}(|\nabla u_2|^{p-2} \nabla u_2) &= \varkappa b(x), \quad \text{in } \Omega, \\ u_2 &= 0, \quad \text{on } \partial\Omega, \end{aligned}$$

where \varkappa is a positive parameter.

It is easy to see that $u_2 \in C^{1,\alpha}(\overline{\Omega})$. Then $\frac{\partial u_2}{\partial \nu} > 0$ on $\partial\Omega$, where ν is the inward unit normal vector. We can also see that $u_2 > 0$ on $\partial(\Omega \setminus \overline{\Omega_\varepsilon})$ when $\varepsilon \in (0, \delta)$ is small enough. Let \varkappa be large enough, we have $u_2 \geq 2C_\#$ on $\partial(\Omega \setminus \overline{\Omega_\varepsilon})$. It means

that $u_2 M^{\frac{1}{p^- - 1}} \geq u$ on $\partial\Omega_\varepsilon$. Define $u_3 = u_2 M^{\frac{1}{p^- - 1}}$. Since $p(x) \equiv p$ for any $x \in \Omega$ with $d(x) \leq \delta$, we have

$$-\operatorname{div}(|\nabla u_3|^{p(x)-2} \nabla u_3) = -\operatorname{div}(|\nabla u_3|^{p-2} \nabla u_3) = M^{\frac{p-1}{p^- - 1}} \kappa b(x) \geq Mb(x) \quad \text{in } \Omega_\varepsilon.$$

Therefore, $u_3 = u_2 M^{\frac{1}{p^- - 1}} \geq u$ on $\overline{\Omega_\varepsilon}$ and $|\nabla u| \leq |\nabla u_3| \leq CM^{\frac{1}{p^- - 1}}$ on $\partial\Omega$. The proof is complete. \square

Proof of Theorem 1.2. At first, we construct a supersolution of (1.1). Denote $k_2 = \psi(k_1\sigma)$. Let ω_2 be the solution of the problem

$$\begin{aligned} -\Delta_{p(x)} \omega_2 &= b(x) k_2^{p^- - 1 - \frac{\varepsilon}{2}}, \quad \text{in } \Omega \setminus \overline{\Omega_\sigma}, \\ \omega_2 &> 0, \quad \text{in } \Omega \setminus \overline{\Omega_\sigma}, \\ \omega_2 &= 0, \quad \text{on } \partial(\Omega \setminus \overline{\Omega_\sigma}). \end{aligned}$$

Define

$$v_2(x) := \begin{cases} \psi(k_1\omega_b(x)), & x \in \Omega_\sigma, \\ \omega_2(x) + \psi(k_1\sigma), & x \in \Omega \setminus \Omega_\sigma. \end{cases} \quad (3.25)$$

For a large enough constant k_1 , we will prove that v_2 is a supersolution of (1.1) in three steps.

Step 1. When k_1 is large enough, we will check that v_2 is a supersolution of (1.1) in Ω_σ , namely,

$$\int_{\Omega} |\nabla v_2|^{p(x)-2} \nabla v_2 \nabla \phi \, dx \geq \int_{\Omega} \lambda b g(v_2) \phi \, dx \geq \int_{\Omega} \lambda f(x, v_2) \phi \, dx,$$

for any $\phi \in C_0^\infty(\Omega_\sigma)$ with $\phi \geq 0$ and $\operatorname{supp} \phi \subset \subset \Omega_\sigma$. As in the proof of Lemma 3.4, we only need to prove that

$$\begin{aligned} &\int_{\Omega} (k_1)^{p(x)} |\nabla \omega_b|^{p(x)} \phi(p(x) - 1) [H(v_2)]^{p(x)-1} \frac{2^\varepsilon - 1}{p^- - 1} \frac{H(v_2)}{v_2} \, dx \\ &\geq \int_{\Omega} (k_1)^{p(x)-1} |\nabla \omega_b|^{p(x)-1} |\nabla p| [\phi \psi'(k_1\omega_b)^{p(x)-1}] |\ln k_1 H(v_2)| \, dx, \end{aligned} \quad (3.26)$$

$$k_1 H(v_2(x)) \geq 1, \quad \forall x \in \Omega_\sigma, \quad (3.27)$$

$$\frac{k_1^{p^- - 1}}{[v_2(x)]^{p^- - 1}} \geq 1, \quad \forall x \in \Omega_\sigma. \quad (3.28)$$

We can see that (3.26) is valid, provided

$$|\nabla \omega_b| \frac{2^\varepsilon - 1}{p^- - 1} k_1 \frac{H(v_2)}{v_2} \geq c_1 \frac{2^\varepsilon - 1}{p^- - 1} k_1 \frac{H(v_2)}{v_2} \geq |\nabla p| |\ln k_1 H(v_2)| \quad \text{in } \Omega_\sigma. \quad (3.29)$$

According to the assumption on g , without loss of generality, we assume that

$$g(s) \geq cs^{-1} \quad \text{for } s \leq 1, \quad \text{and} \quad g(s) = cs^\theta \quad \text{for } s \geq 1,$$

where $\theta = p^- - 1 - \varepsilon$. Thus

$$\hat{g}(s) \geq cs^{-p^-} \quad \text{for } s \leq 1, \quad \text{and} \quad \hat{g}(s) = cs^{\theta+1-p^-} \quad \text{for } s \geq 1,$$

$$H(s) \geq c_1 s^{-\frac{p^-}{p^- - 1}} \quad \text{for } s \leq 1, \quad \text{and} \quad H(s) = c_2 s^{\frac{\theta+1-p^-}{p^- - 1}} \quad \text{for } s \geq 2,$$

$$\eta(s) \leq c_3 s^{1+\frac{p^-}{p^- - 1}} \quad \text{for } s \leq 1, \quad \text{and} \quad c_4 s^{2-\frac{\theta}{p^- - 1}} \leq \eta(s) \leq c_5 s^{2-\frac{\theta}{p^- - 1}} \quad \text{for } s \geq 3.$$

Then $\psi(s)$ satisfies

$$c_7 s^{\frac{1}{2-\frac{\theta}{p^-}-1}} \leq \psi(s) \leq c_8 s^{\frac{1}{2-\frac{\theta}{p^-}-1}} \quad \text{for } s \geq 3.$$

Let $s_0 \geq 3$ such that $\eta(s_0) \geq 3$. Denote

$$\Omega_\sigma^+ = \{x \in \Omega_\sigma : k_1 \omega_b(x) \geq \eta(s_0)\}, \quad \Omega_\sigma^- = \{x \in \Omega_\sigma : k_1 \omega_b(x) < \eta(s_0)\}.$$

Here we note that $v_2 = \psi(k_1 \omega_b)$ on $\overline{\Omega_\sigma^-}$. Since ψ is strictly increasing, we have $k_1 \omega_b \geq \eta(s_0)$ if and only if $v_2 = \psi(k_1 \omega_b) \geq \psi(\eta(s_0)) = s_0 \geq 3$. When $v_2 \geq s_0$, we have

$$\begin{aligned} c_7(k_1 \omega_b)^{\frac{1}{2-\frac{\theta}{p^-}-1}} &\leq v_2 \leq c_8(k_1 \omega_b)^{\frac{1}{2-\frac{\theta}{p^-}-1}} \quad \text{on } \Omega_\sigma^+, \\ c_9(k_1 \omega_b)^{\frac{1}{2-\frac{\theta}{p^-}-1} \frac{\theta+1-p^-}{p^-}} &\leq H(v_2) \leq c_{10}(k_1 \omega_b)^{\frac{1}{2-\frac{\theta}{p^-}-1} \frac{\theta+1-p^-}{p^-}} \quad \text{on } \Omega_\sigma^+, \\ |\nabla \omega_b| \frac{2^\varepsilon - 1}{p^- - 1} k_1 \frac{H(v_2)}{v_2} &\geq c_{11} k_1 (k_1 \omega_b)^{\frac{1}{2-\frac{\theta}{p^-}-1} (\frac{\theta+1-p^-}{p^-} - 1)} = \frac{c_{11}}{\omega_b} \geq \frac{c_{11}}{\sigma} \quad \text{on } \Omega_\sigma^+, \\ |\nabla p| \ln k_1 H(v_2) &\leq |\nabla p| (\ln k_1 + |\ln H(v_2)|) \\ &\leq |\nabla p| (\ln k_1 + c_{12} |\ln k_1 \omega_b|) \leq c_{13} \ln k_1 \quad \text{on } \Omega_\sigma^+. \end{aligned}$$

Denoting $\sigma = \frac{c_{11}}{c_{13} \ln k_1}$, we obtain

$$|\nabla \omega_b| \frac{2^\varepsilon - 1}{p^- - 1} k_1 \frac{H(v_2)}{v_2} \geq c_1 \frac{2^\varepsilon - 1}{p^- - 1} k_1 \frac{H(v_2)}{v_2} \geq |\nabla p| \ln k_1 H(v_2) \quad \text{on } \Omega_\sigma^+.$$

Since ψ is strictly increasing, we have $k_1 \omega_b \leq \eta(s_0)$ if and only if $v_2 = \psi(k_1 \omega_b) \leq s_0$. Note that $H(v_2)$ is decreasing. It follows that $H(v_2) \geq H(s_0)$ on Ω_σ^- . Thus (3.29) is valid when k_1 is large enough. Thus (3.29) is valid, and then (3.26) is valid.

Obviously,

$$\begin{aligned} k_1 H(v_2(x)) &\geq k_1 H(\psi(k_1 \sigma)) \\ &\geq k_1 c_9(k_1 \frac{c_{11}}{c_{13} \ln k_1})^{\frac{1}{2-\frac{\theta}{p^-}-1} \frac{\theta+1-p^-}{p^-}} \\ &= c_9(k_1)^{\frac{1}{2-\frac{\theta}{p^-}-1}} (\frac{c_{11}}{c_{13} \ln k_1})^{\frac{1}{2-\frac{\theta}{p^-}-1} \frac{\theta+1-p^-}{p^-}} \rightarrow +\infty, \end{aligned}$$

for all $x \in \Omega_\sigma$ as $k_1 \rightarrow +\infty$. Thus (3.27) is valid.

Note that $\frac{\theta}{p^- - 1} < 1$. Then by the above computation,

$$v_2 \leq c_8(k_1 \omega_b)^{\frac{1}{2-\frac{\theta}{p^-}-1}} \leq c_8(k_1 \sigma)^{\frac{1}{2-\frac{\theta}{p^-}-1}} \leq k_1$$

as k_1 is large enough. Thus (3.28) is valid.

Step 2. We will check that v_2 is a supersolution of (1.1) in $\Omega \setminus \overline{\Omega_\sigma^-}$ when k_1 is large enough; i.e.,

$$\int_{\Omega} |\nabla v_2|^{p(x)-2} \nabla v_2 \nabla \phi \, dx \geq \int_{\Omega} \lambda b g(v_2) \phi \, dx \geq \int_{\Omega} \lambda f(x, v_2) \phi \, dx,$$

for all $\phi \in C_0^\infty(\Omega \setminus \overline{\Omega_\sigma^-})$, $\phi \geq 0$. By the definition of ω_2 and Lemmas 3.5 and 3.6, we have

$$\omega_2 \leq C_1(k_2)^{\frac{p^- - 1 - \frac{\varepsilon}{2}}{p^- - 1}}, \quad |\nabla \omega_2| \leq C_2(k_2)^{\frac{p^- - 1 - \frac{\varepsilon}{2}}{p^- - 1}}.$$

Since $v_2 = \omega_2 + \psi(k_1\sigma)$ in $\Omega \setminus \overline{\Omega_\sigma}$, we have

$$c_7(k_1\sigma)^{\frac{1}{2-\frac{\theta}{p^-}-1}} \leq \psi(k_1\sigma) \leq v_2 \leq C_1(k_2)^{\frac{p^- - 1 - \frac{\epsilon}{2}}{p^- - 1}} + \psi(k_1\sigma) \leq (C_1 + 1)\psi(k_1\sigma).$$

Since $k_1\sigma = \frac{c_{11}k_1}{c_{13}\ln k_1}$ is large enough (as long as k_1 is large enough) and $v_2(\cdot)$ is large enough in $\Omega \setminus \overline{\Omega_\sigma}$, the assumption (ii) of Theorem 1.2 implies that $\frac{g(v_2(x))}{[v_2(x)]^{p^- - 1 - \frac{\epsilon}{2}}}$ is small enough. Therefore, we see that

$$\frac{1}{\lambda} > \frac{g(v_2(x))}{[v_2(x)]^{p^- - 1 - \frac{\epsilon}{2}}}.$$

Note that $k_2 = \psi(k_1\sigma)$. We have

$$v_2(x)^{p^- - 1 - \frac{\epsilon}{2}} \leq [C_1(k_2)^{\frac{p^- - 1 - \frac{\epsilon}{2}}{p^- - 1}} + \psi(k_1\sigma)]^{p^- - 1 - \frac{\epsilon}{2}} \leq C_3 k_2^{p^- - 1 - \frac{\epsilon}{2}}, \quad \forall x \in \Omega \setminus \overline{\Omega_\sigma},$$

which implies

$$(k_2)^{p^- - 1 - \frac{\epsilon}{2}} \geq C_3 \lambda (k_2)^{p^- - 1 - \frac{\epsilon}{2}} \frac{g(v_2(x))}{[v_2(x)]^{p^- - 1 - \frac{\epsilon}{2}}} \geq \lambda g(v_2(x)), \quad \forall x \in \Omega \setminus \overline{\Omega_\sigma}.$$

We can see that v_2 is a supersolution of (1.1) in $\Omega \setminus \overline{\Omega_\sigma}$; i.e., for any $\phi \in C_0^\infty(\Omega \setminus \overline{\Omega_\sigma})$ with $\phi \geq 0$, we have

$$\begin{aligned} \int_{\Omega} |\nabla v_2|^{p(x)-2} \nabla v_2 \nabla \phi \, dx &= \int_{\Omega} b k_2^{p^- - 1 - \frac{\epsilon}{2}} \phi \, dx \\ &\geq \int_{\Omega} \lambda b C_3 k_2^{p^- - 1 - \frac{\epsilon}{2}} \frac{g(v)}{v_2^{p^- - 1 - \frac{\epsilon}{2}}} \phi \, dx \\ &\geq \int_{\Omega} \lambda f(x, v_2) \phi \, dx. \end{aligned}$$

Step 3. When k_1 is large enough, we will prove that v_2 is a supersolution of (1.1) in Ω . When $\omega_b(x) = \sigma$, it is easy to check that

$$\begin{aligned} k_1 \psi'(k_1 \omega_b) &= k_1 H(v_2) \\ &\geq k_1 c_9 (k_1 \omega_b)^{\frac{1}{2-\frac{\theta}{p^-}-1} \frac{\theta+1-p^-}{p^- - 1}} \\ &= c_9 (k_1)^{\frac{1}{2-\frac{\theta}{p^-}-1}} \sigma^{\frac{1}{2-\frac{\theta}{p^-}-1} \frac{\theta+1-p^-}{p^- - 1}} \\ &= c_9 (k_1)^{\frac{1}{2-\frac{\theta}{p^-}-1}} \left(\frac{c_{11}}{c_{13} \ln k_1} \right)^{\frac{1}{2-\frac{\theta}{p^-}-1} \frac{\theta+1-p^-}{p^- - 1}}. \end{aligned}$$

Then

$$|\nabla \omega_2| \leq C(k_2)^{\frac{p^- - 1 - \frac{\epsilon}{2}}{p^- - 1}} \leq C \left(\frac{c_{11}k_1}{c_{13} \ln k_1} \right)^{\frac{1}{2-\frac{\theta}{p^-}-1} \frac{p^- - 1 - \frac{\epsilon}{2}}{p^- - 1}} < |\nabla \psi(k_1 \omega_b)|$$

as $k_1 \rightarrow +\infty$. Thus we know that

$$(k_1 \psi'(k_1 \omega_b(x)) |\nabla \omega_b(x)|)^{p(x)-1} - |\nabla \omega_2(x)|^{p(x)-1} > 0, \quad \text{when } \omega_b(x) = \sigma.$$

Therefore, when $\sigma = c_{11}/(c_{13} \ln k_1)$ and k_1 is large enough, similar argument as to the step 3 of the proof of Lemma 3.4 implies v_2 is a supersolution of (1.1).

It is easy to see that $\mu\phi$ defined in the proof of Theorem 1.1 is a subsolution of (1.1) and $\mu\phi \leq v_2$ when μ is small enough. By Lemma 2.7, we can get the existence of a solution to (1.1). The proof is complete. \square

Proof of Theorem 1.3. At first, similar to the proof of Theorem 1.2, we will prove that v_2 defined by (3.25) is also a supersolution of (1.1) for a large enough constant k_1 .

Similar to the proof of Theorem 1.2, we consider the solution ω_2 of the problem

$$\begin{aligned} -\Delta_{p(x)}\omega_2 &= b(x)k_2^{p^- - 1 - \frac{\varepsilon}{2}}, \quad \text{in } \Omega \setminus \overline{\Omega_\sigma}, \\ \omega_2(x) &> 0, \quad \text{in } \Omega \setminus \overline{\Omega_\sigma}, \\ \omega_2(x) &= 0, \quad \text{on } \partial(\Omega \setminus \overline{\Omega_\sigma}), \end{aligned} \quad (3.30)$$

where $k_2 = \psi(k_1\sigma)$. We have

$$\omega_2 \leq C_1(k_2)^{\frac{p^- - 1 - \frac{\varepsilon}{2}}{p^- - 1}}.$$

Next we only need to prove that

$$|\nabla\omega_2| \leq C_2(k_2)^{\frac{p^- - 1 - \frac{\varepsilon}{2}}{p^- - 1}}.$$

Now we consider $(\gamma k_2)^{\frac{p^- - 1 - \frac{\varepsilon}{2}}{p^- - 1}}\omega_b$, where $\gamma \geq 1$ is a constant. Here we note that $\nabla\omega_b \cdot \nu = |\nabla\omega_b|$ on $\partial(\Omega \setminus \overline{\Omega_\sigma})$ and $\nabla p \cdot \nu < 0$ on $\partial(\Omega \setminus \overline{\Omega_\sigma})$, where ν is the inward unit normal vector on $\partial(\Omega \setminus \overline{\Omega_\sigma})$. There exists a small enough positive constant $\delta > 0$ such that $\nabla\omega_b \nabla p < 0$ in $(\Omega \setminus \overline{\Omega_\sigma})_\delta^\# := \{x \in \Omega \setminus \overline{\Omega_\sigma} : d(x, \partial(\Omega \setminus \overline{\Omega_\sigma})) < \delta\}$. By computations it follows that

$$\begin{aligned} & -\Delta_{p(x)}(\gamma k_2)^{\frac{p^- - 1 - \frac{\varepsilon}{2}}{p^- - 1}}\omega_b \\ &= (\gamma k_2)^{\frac{p^- - 1 - \frac{\varepsilon}{2}}{p^- - 1}(p(x)-1)}(-\Delta_{p(x)}\omega_b - \frac{p^- - 1 - \frac{\varepsilon}{2}}{p^- - 1}|\nabla\omega_b|^{p(x)-2}\nabla\omega_b \nabla p \ln \gamma k_2) \\ &\geq (\gamma k_2)^{\frac{p^- - 1 - \frac{\varepsilon}{2}}{p^- - 1}(p(x)-1)}(-\Delta_{p(x)}\omega_b) \\ &\geq b(\gamma k_2)^{p^- - 1 - \frac{\varepsilon}{2}} \text{ in } (\Omega \setminus \overline{\Omega_\sigma})_\delta^\#. \end{aligned}$$

Since ω_b is positive and continuous, there exists a large enough positive γ such that $\gamma^{\frac{p^- - 1 - \frac{\varepsilon}{2}}{p^- - 1}}\omega_b > 2C_1$ for $d(x, \partial(\Omega \setminus \overline{\Omega_\sigma})) = \delta$. Therefore $(\gamma k_2)^{\frac{p^- - 1 - \frac{\varepsilon}{2}}{p^- - 1}}\omega_b$ is a supersolution of (3.30) in $(\Omega \setminus \overline{\Omega_\sigma})_\delta^\#$. By the comparison principle, we have $(\gamma k_2)^{\frac{p^- - 1 - \frac{\varepsilon}{2}}{p^- - 1}}\omega_b \geq \omega_2$ in $(\Omega \setminus \overline{\Omega_\sigma})_\delta^\#$, and then

$$|\nabla\omega_2| \leq |(\gamma k_2)^{\frac{p^- - 1 - \frac{\varepsilon}{2}}{p^- - 1}}\nabla\omega_b| \leq C_2(k_2)^{\frac{p^- - 1 - \frac{\varepsilon}{2}}{p^- - 1}} \text{ on } \partial(\Omega \setminus \overline{\Omega_\sigma}).$$

Note that

$$\max_{x \in \Omega \setminus \overline{\Omega_\sigma}} \omega_2(x) \leq C_3(k_2)^{\frac{p^- - 1 - \frac{\varepsilon}{2}}{p^- - 1}}$$

and $k_2 = \psi(k_1\sigma)$. Since $v_2 = \omega_2 + \psi(k_1\sigma)$ in $\Omega \setminus \overline{\Omega_\sigma}$, we have

$$\begin{aligned} c_7(k_1\sigma)^{\frac{1}{2 - \frac{\theta}{p^- - 1}}} &\leq \psi(k_1\sigma) \leq v_2 \\ &\leq \max_{x \in \Omega \setminus \overline{\Omega_\sigma}} \omega_2(x) + \psi(k_1\sigma) \end{aligned}$$

$$\leq C_3(k_2)^{\frac{p^- - 1 - \frac{\varepsilon}{2}}{p^- - 1}} + \psi(k_1\sigma) \leq C_4\psi(k_1\sigma).$$

As in the proof of Theorem 1.2, we can see that the v_2 defined in the proof of Theorem 1.2 is a supersolution of (1.1).

It is easy to see that $\mu\phi$ defined in the proof of Theorem 1.1 is a subsolution of (1.1), and $\mu\phi \leq v_2$ when μ is small enough. By Lemma 2.7, we obtain the existence of solution of (1.1). The proof is complete. \square

Proof of Theorem 1.4. The proof is similar to that of Theorem 1.2. We will prove that v_2 defined in (3.25) is a supersolution of (1.1) for a large enough constant k_1 .

Since (1.1) is radial, we may assume the both solutions $\omega_b(\cdot)$ and $\omega_2(\cdot)$ are radial. We only need to prove that v_2 defined in (3.25) is also a supersolution of (1.1) for a large enough constant k_1 . Since ω_2 is radial, it is easy to see that

$$\omega_2 \leq C_1(k_2)^{\frac{p^- - 1 - \frac{\varepsilon}{2}}{p^- - 1}}, |\nabla \omega_2| \leq C_2(k_2)^{\frac{p^- - 1 - \frac{\varepsilon}{2}}{p^- - 1}}.$$

Similar to the proof of Theorem 1.2, we can see that the v_2 defined in (3.25) is a supersolution of (1.1). The proof is complete. \square

Acknowledgments. The authors would like to thank Professor Julio G. Dix for his suggestions, and the anonymous referees for their valuable comments.

This research was partially supported by the National Natural Science Foundation of China (11326161 and 10971087) and the key projects of Science and Technology Research of the Henan Education Department (14A110011).

REFERENCES

- [1] E. Acerbi, G. Mingione; Regularity results for a class of functionals with nonstandard growth, *Arch. Ration. Mech. Anal.* 156 (2001), 121–140.
- [2] N. Benouhiba; On the eigenvalues of weighted $p(x)$ -Laplacian on \mathbb{R}^N , *Nonlinear Anal.*, 74(2011)235-243.
- [3] F. Cammaroto, L. Vilasi; On a perturbed $p(x)$ -Laplacian problem in bounded and unbounded domains, *J. Math. Anal. Appl.*, 402 (2013), 71-83.
- [4] K.C. Chang; *Critical point theory and applications*, Shanghai Scientific and Technology press, Shanghai, 1986.
- [5] Y. Chen, S. Levine, M. Rao; Variable exponent, linear growth functionals in image restoration, *SIAM J. Appl. Math.*, **66** (2006), No.4, 1383–1406.
- [6] L. Diening, P. Harjulehto, P. Hästö, M. Růžička; *Lebesgue and Sobolev Spaces with Variable Exponents*, Lecture Notes in Mathematics 2017, Springer-Verlag, Berlin, 2011.
- [7] X. L. Fan; Global $C^{1,\alpha}$ regularity for variable exponent elliptic equations in divergence form, *J. Differ. Equations* 235 (2007) 397-417.
- [8] X. L. Fan; Eigenvalue of the $p(x)$ -Laplacian Numann problems, *Nonlinear Anal.*, 67 (2007), 2982-2992.
- [9] X. L. Fan; On the sub-super solution method for $p(x)$ -Laplacian equations, *J. Math. Anal. Appl.* 330 (2007), 665-682.
- [10] X. L. Fan, D. Zhao; A class of De Giorgi type and Hölder continuity, *Nonlinear Anal.*, 36 (1999), 295-318.
- [11] X. L. Fan, D. Zhao; The quasi-minimizer of integral functionals with $m(x)$ growth conditions, *Nonlinear Anal.*, 39 (2000) 807-816.
- [12] X. L. Fan, Q. H. Zhang; Existence of solutions for $p(x)$ -Laplacian Dirichlet problem, *Nonlinear Anal.*, 52(2003)1843-1852.
- [13] X. L. Fan, Q. H. Zhang, D. Zhao; Eigenvalues of $p(x)$ -Laplacian Dirichlet problem, *J. Math. Anal. Appl.* 302(2005)306-317.
- [14] X. L. Fan, D. Zhao; On spaces $L^{p(x)}(\Omega)$ and $W^{m,p(x)}(\Omega)$, *J. Math. Anal. Appl.* 263 (2001) 424-446.

- [15] J. V. A. Gonçalves, M. C. Rezende, C. A. Santos; Positive solutions for a mixed and singular quasilinear problem, *Nonlinear Anal.*, 74(2011)132-140.
- [16] Z. C. Guo, Q. Liu, J. B. Sun, B. Y. Wu; Reaction-diffusion systems with $p(x)$ -growth for image denoising, *Nonlinear Anal-Real.*, 12 (2011), 2904-2918.
- [17] Z. C. Guo, J. B. Sun, D. Z. Zhang, B. Y. Wu; Adaptive perona-malik model based on the variable exponent for image denoising, *IEEE T. Image Process*, 21 (2012), 958-967.
- [18] C. M. Guo, C. B. Zhai, R. P. Song; An existence and uniqueness result for the singular Lane-Emden-Fowler equation, *Nonlinear Anal.*, 72 (2010), 1275-1279.
- [19] T. C. Halsey; *Electrorheological fluids*, Science 258 (5083) (1992), 761-766.
- [20] A. El Hamidi; Existence results to elliptic systems with nonstandard growth conditions, *J. Math. Anal. Appl.*, 300 (2004), 30-42.
- [21] P. Harjulehto, P. Hästö, M. Koskenoja, S. Varonen; The Dirichlet energy integral and variable exponent Sobolev spaces with zero boundary values, *Potential Anal.* 25 (2006), 205-222.
- [22] P. Harjulehto, P. Hästö, V. Latvala; Harnack's inequality for $p(\cdot)$ -harmonic functions with unbounded exponent p , *J. Math. Anal. Appl.*, 352 (2009), 345-359.
- [23] P. Harjulehto, P. Hästö, V. Latvala, O. Toivanen; Critical variable exponent functionals in image restoration, *Appl. Math. Lett.*, 26 (2013), 56-60.
- [24] P. Harjulehto, P. Hästö, Ú. V. Lê, M. Nuortio; Overview of differential equations with non-standard growth, *Nonlinear Anal.*, 72 (2010), 4551-4574.
- [25] P. Harjulehto, P. Hästö, V. Latvala; Minimizers of the variable exponent, non-uniformly convex Dirichlet energy, *J. Math. Pure. Appl.*, 89 (2008) (2), pp. 174-197.
- [26] O. Kováčik, J. Rákosník; On spaces $L^{p(x)}(\Omega)$ and $W^{k,p(x)}(\Omega)$, *Czech. Math. J.*, 41 (1991), 592-618.
- [27] O. A. Ladyzenskaja, N. N. Ural'tzeva; *Linear and Quasilinear Elliptic Equations*, Academic Press, New York, 1968.
- [28] F. Li, Z. B. Li, L. Pi; Variable exponent functionals in image restoration, *Appl. Math. Comput.*, 216 (2010), 870-882.
- [29] J. J. Liu; Positive solutions of the $p(x)$ -Laplace equation with singular nonlinearity, *Nonlinear Anal.*, 72 (2010), 4428-4437.
- [30] T. Lukkari; Singular solutions of elliptic equations with nonstandard growth, *Math. Nachr.*, 282 (2009), 1770-1787.
- [31] M. Mihăilescu, V. Rădulescu; Continuous spectrum for a class of nonhomogeneous differential operators, *Manuscripta Math.* 125 (2008), 157-167.
- [32] M. Mihăilescu, V. Rădulescu; On a nonhomogeneous quasilinear eigenvalue problem in Sobolev spaces with variable exponent, *Proc. Amer. Math. Soc.* 135 (2007), 2929-2937.
- [33] M. Mihăilescu, V. Rădulescu, D. Stancu-Dumitru; A Caffarelli-Kohn-Nirenberg-type inequality with variable exponent and applications to PDE's, *Complex Var. Elliptic Equa.*, 56 (2011), 659-669.
- [34] M. Mihăilescu, G. Moroşanu, D. Stancu-Dumitru; Equations involving a variable exponent Grushin-type operator, *Nonlinearity*, 24 (2011), 2663-2680.
- [35] M. Mihăilescu, V. Rădulescu; Concentration phenomena in nonlinear eigenvalue problems with variable exponents and sign-changing potential, *J. Anal. Math.*, 111 (2010), 267-287.
- [36] M. Mihăilescu, V. Rădulescu, D. Repovš; On a non-homogeneous eigenvalue problem involving a potential: An Orlicz-Sobolev space setting, *J. Math. Pures Appl.*, 93 (2010) (2), 132-148.
- [37] A. Mohammed; Positive solutions of the p -Laplace equation with singular nonlinearity, *J. Math. Anal. Appl.* 352 (2009), 234-245.
- [38] K. Perera, E. A. B. Silva; On singular p -Laplacian problems, *Differential Integral Equations*, 20 (2007), 105-120.
- [39] P. Pucci, Q.H. Zhang; Existence of entire solutions for a class of variable exponent elliptic equations, *J. Differ. Equations*, 257 (2014), 1529-1566.
- [40] M. Qing, Z. Yang; Quasilinear elliptic equation involving singular nonlinearities, *Int. J. Comput. Math.* (2009), 1-11.
- [41] K. R. Rajagopal, M. Růžička; Mathematical modeling of electrorheological fluids, *Contin. Mech. Thermodyn.* 13 (2001), 59-78.
- [42] M. Růžička; *Electrorheological fluids: modeling and mathematical theory*, Lecture Notes in Math 1748, Springer-Verlag, Berlin, 2000.
- [43] Q. H. Zhang; A strong maximum principle for differential equations with nonstandard $p(x)$ -growth conditions, *J. Math. Anal. Appl.* 312(2005), No.1, 24-32.

- [44] Q. H. Zhang; Existence and asymptotic behavior of positive solutions to $p(x)$ -Laplacian equations with singular nonlinearities, *J. Ineq. Appl.*, 2007 (2007), Article ID 19349, 9 pages.
- [45] Q. H. Zhang; Existence, nonexistence and asymptotic behavior of boundary blow-up solutions to $p(x)$ -Laplacian problems with singular coefficient, *Nonlinear Anal.*, 74 (2011), 2045–2061.
- [46] V. V. Zhikov; Averaging of functionals of the calculus of variations and elasticity theory, *Math. USSR. Izv.* **29** (1987), 33–36.

JINGJING LIU

COLLEGE OF MATHEMATICS AND INFORMATION SCIENCE, ZHENGZHOU UNIVERSITY OF LIGHT INDUSTRY, ZHENGZHOU, HENAN 450002, CHINA

E-mail address: `jingjing830306@163.com`

QIHU ZHANG (CORRESPONDING AUTHOR)

COLLEGE OF MATHEMATICS AND INFORMATION SCIENCE, ZHENGZHOU UNIVERSITY OF LIGHT INDUSTRY, ZHENGZHOU, HENAN 450002, CHINA

E-mail address: `zhangqihu@yahoo.com`, `zhangqh1999@yahoo.com.cn`

CHUNSHAN ZHAO

DEPARTMENT OF MATHEMATICAL SCIENCES, GEORGIA SOUTHERN UNIVERSITY, STATESBORO, GA 30460, USA

E-mail address: `czhao@GeorgiaSouthern.edu`